

Well Visit & Shot Schedule

Christopher Bassett, MD
Cowles Clinic Pediatrics (706) 454-1210

This is our usual schedule of well child exams and immunizations. This schedule does change often, when there are national shortages of certain vaccines, or when new well-care recommendations are made.

Newborn visit at 3-5 days of life

2 week well visit

- HepB #1 vaccine given if not given at birth

1 month weight check

- For premature or smaller babies

2 month well visit

- DTAP #1
- IPV #1
- Hib #1
- Prevnar #1
- Rota #1

4 month well visit

- DTAP #2
- IPV #2
- Hib #2
- Prevnar #2
- Rota #2

6 month well visit

- DTAP #3
- IPV #3
- Prevnar #3
- Rota #3

9 month well visit

- HepB #2
- Catch-up shots if any are missing
- Finger prick to check for anemia

12 month well visit

- MMR #1
- VVX #1
- HepA #1

15 month well visit

- DTaP #4
- Hib #3
- Prevnar #4

18 month well visit

- HepA #2
- HepB #3

2 year and 3 year well visits

- Catch-up shots if any are missing

4 year well visit

- DTAP #5
- IPV #4
- MMR #2
- VVX #2

5 year well visit

- Catch-up shots if any are missing
- Urinalysis and finger prick

Well visits at 6, 7, 8, 9, and 10 years.**11-12 year well visit**

- **Menactra**
- **TdaP**
- **Gardasil (for girls, repeated two more times)**

Immunization abbreviations / descriptions:

DTaP: Diphtheria, Tetanus and Pertussis (Whooping cough) prevention - need 5 total

Gardasil: Gardasil shot, prevents genital warts and cervical cancer in girls - need 3

HepB: Hepatitis B, prevents liver infection by hepatitis B - need 3

HepA: Hepatitis A, prevents liver infection by hepatitis A - need 2

Hib: Haemophilus Influenzae B, prevents brain/blood infections by this bacterium - need 3

IPV: Polio, prevents poliomyelitis - need 4

Menactra: Neisseria meningitidis, prevents brain/blood infections by this bacterium - need 1

MMR: Measles, Mumps, and Rubella prevention - need 2

Prevnar: Streptococcus pneumoniae, prevents brain/blood infections by this bacterium - need 4

Rota: Rotavirus prevention - need 3

TdaP: Tetanus and Pertussis (whooping cough) booster - need 1

VVX: Varicella or "Chicken Pox" prevention - need 2